

High-Resolution Bio-Optical Property Fields in the Northern Gulf of Mexico: Five-Year Ocean Color Time Series




***Richard W. Gould, Jr.
Jason Jolliff
Sherwin D. Ladner***

***Naval Research Laboratory
Bio-Optical/Physical Processes and Remote Sensing Section,
Code 7331
Oceanography Division
Stennis Space Center, Mississippi***

Objectives

- Establish baseline bio-optical properties for coastal habitats in the Northern Gulf of Mexico (5-year satellite climatology).
- Communicate this information to environmental resource managers to aid decision making (dredging operations,




bio-Optical Products - MODIS Aqua, high-resolution (250m)


for each bay and sub-region; weekly, monthly products):

Total Suspended Solids (TSS) □ Particulate Inorganic Matter (PIM)
Particulate Organic Matter (POM) □ Diffuse Attenuation Coefficients at 488nm (K_d)
Euphotic Depth (Zeu)


Mobile Bay, AL




Pensacola Bay, FL



Choctawhatchee Bay, FL



Saint Andrew Bay, FL



Hyperspectral Imager for the Coastal Ocean (HICO)

NRL controls HICO operations HICO Processing

- *Target selection*
- *Target acquisition*

HICO Specifications

- *100m spatial resolution*
- *87 spectral channels (400-900 μ m)*
- *5.7 μ m spectral resolution*

Optical Products

Chlorophyll

Absorption coefficient

(phyto, CDOM, sediment/detritus)

Backscattering coefficient

Diffuse attenuation coefficient

Beam attenuation coefficient

Euphotic depth

TSS (organic, inorganic)


Optical water mass classification

- *Convolved to “MODIS-like” bands*
(multispectral)
- *Full hyperspectral*
- *Consistent with processing for other ocean color sensors (SeaWiFS, MODIS, MERIS, VIIRS)*




9 December 2009
Mobile Bay
(descending pass)

True Color




Mississippi-Alabama Shelf

9 December 2009
Mobile Bay
(descending pass)




9 December 2009
Mobile Bay
(descending pass)

TSS



Mobile Bay, AL

MODIS time-series



HICO (single image)



- *not exactly the same areas covered by the sub-regions.*
- *weekly mean vs. individual scene.*

MODIS (weekly mean, 12/3-12/10/2009) vs. HICO

Region	Property	MODIS time-series	HICO	% Difference (HICO - MODIS)
Plume	TSS	4.0	4.7	17.5
	PIM	3.1	3.9	25.8
	POM	1.0	0.7	30.0
	Zeu	11.0	9.0	18.2
	K _d (488)	0.63	0.61	3.2
Lower Bay	TSS	9.5	6.5	31.6
	PIM	8.1	5.6	30.9
	POM	1.4	0.8	42.8
	Zeu	4.5	6.6	46.7
	K _d (488)	1.68	0.96	42.8
Upper Bay	TSS	11.8	17.7	50.0
	PIM	10.4	16.7	60.6
	POM	1.4	1.0	28.6
	Zeu	6.1	3.1	49.2
	K (488)	1.89	2.14	13.2

generally
increasing
difference
s




11 September 2011
Pensacola Bay
(ascending pass)

True Color

Dauphin Island

Mississippi-Alabama St.

11 September 2011
Pensacola Bay
(ascending pass)



11 September 2011
Pensacola Bay
(ascending pass)

SPM TSS

Pensacola

Santa Rosa Island

Dauphin Island


Island

Mississippi-Alabama Shelf

Time-Series Analysis - MODIS High-Resolution (250m) Bio-Optical

Monthly Composites (2005-2009), $K_d(488)$



Products



Time-Series Analysis - MODIS High-Resolution (250m) Bio-Optical

Monthly Composites (2005-2009), Z_{eu}



Products



Time-Series Analysis - MODIS High-Resolution (250m) Bio-Optical

Monthly Composites (2005-2009), PIM



Products



Time-Series Analysis - MODIS High-Resolution (250m) Bio-Optical

Monthly Composites (2005-2009), POM



Products



Time-Series Analysis - MODIS High-Resolution (250m) Bio-Optical



Monthly Composites (2005-2009), TSS

Products






Time-Series Analysis - Within Bays

5-Year Monthly Averages (2005-2009), Mobile Bay





Time-Series Analysis - Within Bays

5-Year Monthly Averages (2005-2009), Mobile Bay






Time-Series Analysis - Within Bays

5-Year Monthly Averages (2005-2009), Pensacola Bay





Time-Series Analysis - Within Bays

5-Year Monthly Averages (2005-2009), Pensacola Bay






Time-Series Analysis - Within Bays

5-Year Monthly Averages (2005-2009), Choctawhatchee Bay





Time-Series Analysis - Within Bays

5-Year Monthly Averages (2005-2009), Choctawhatchee Bay






Time-Series Analysis - Within Bays

5-Year Monthly Averages (2005-2009), St. Andrew Bay





Time-Series Analysis - Within Bays

5-Year Monthly Averages (2005-2009), St. Andrew Bay



Time-Series Analysis - Within Bays




Weekly Averages For 5 Years (1/1/05-12/31/09),
Mobile Bay



Time-Series Analysis - Within Bays



Weekly Averages For 5 Years (1/1/05-12/31/09),

Mobile Bay



Time-Series Analysis - Within Bays




Weekly Averages For 5 Years (1/1/05-12/31/09),
Mobile Bay



Time-Series Analysis - Within Bays



Weekly Averages For 5 Years (1/1/05-12/31/09),

Mobile Bay



Time-Series Analysis - Within Bays




Weekly Averages For 5 Years (1/1/05-12/31/09),
Pensacola Bay



Time-Series Analysis - Within Bays



Weekly Averages For 5 Years (1/1/05-12/31/09),

Bonacola Bay





Time-Series Analysis - Within Bays

Weekly Averages For 5 Years (1/1/05-12/31/09),
Pensacola Bay



Time-Series Analysis - Within Bays




Weekly Averages For 5 Years (1/1/05-12/31/09),
Pensacola Bay



Time-Series Analysis - Within Bays



Weekly Averages For 5 Years (1/1/05-12/31/09),

Pensacola Bay



Time-Series Analysis - Within Bays




Weekly Averages For 5 Years (1/1/05-12/31/09),
Choctawhatchee Bay



Time-Series Analysis - Within Bays



Weekly Averages For 5 Years (1/1/05-12/31/09),

Choctawhatchee Bay



Time-Series Analysis - Within Bays




Weekly Averages For 5 Years (1/1/05-12/31/09),
Choctawhatchee Bay



Time-Series Analysis - Within Bays



Weekly Averages For 5 Years (1/1/05-12/31/09),

Choctawhatchee Bay



Time-Series Analysis - Within Bays




Weekly Averages For 5 Years (1/1/05-12/31/09), St.
Andrew Bay



Time-Series Analysis - Within Bays



Weekly Averages For 5 Years (1/1/05-12/31/09), St.

Andrew Bay



Time-Series Analysis - Within Bays





Weekly Averages For 5 Years (1/1/05-12/31/09), St.
Andrew Bay



Time-Series Analysis - Within Bays

Weekly Averages For 5 Years (1/1/05-12/31/09), St.




Andrew Bay



Time-Series Analysis - Between Bays



Weekly Averages For 5 Years (1/1/05-12/31/09),

Plumes






Time-Series Analysis - Between Bays

Weekly Averages For 5 Years (1/1/05-12/31/09),
Plumes





Time-Series Analysis - Between Bays

Weekly Averages For 5 Years (1/1/05-12/31/09), Bays






Time-Series Analysis - Between Bays




Weekly Averages For 5 Years (1/1/05-12/31/09), Bays






Bio-Optical Property Relationships - Weekly Averages For 5 Years (1/1/05-12/31/09), Within Bays






Bio-Optical Property Relationships - Weekly Averages For 5 Years (1/1/05-12/31/09), Within Bays






Bio-Optical Property Relationships - Weekly Averages For 5 Years (1/1/05-12/31/09), Choctawhatchee Bay



Bio-Optical Property Relationships - Weekly Averages For 5 Years (1/1/05-12/31/09), St. Andrew Bay




Bio-Optical Property Relationships - Weekly Averages For 5 Years (1/1/05-12/31/09), All Regions In Each Bay



Bio-Optical Property

Turbidity (NTU) vs. Total Suspended Sediments (TSS)

Relationships



- Multiple cruises (6)
- Mississippi Bight, Mobile Bay samples
- Gravimetric TSS, Turner Aquafluor turbidity (NTU)



United States Naval Research Laboratory
Bio-Optical/Physical Processes and Remote Sensing Section

[HOME](#)[BROWSE IMAGERY](#)[EXERCISE SUPPORT](#)[EXTERNAL LINKS](#)[PROJECTS](#)[CURRENT PROJECTS](#)[PUBLICATIONS](#)

CLICK HERE: High-Resolution Subsurface Physical and Optical Property Fields in the Gulf of Mexico.
Establishing Baselines and Assessment Tools for Resource Managers

[Home Page Navigation](#)[Section Info](#)[Regions of Interest Info](#)[Browse Imagery Info](#)[HICO Archive Target Search](#)[Mobile Image Tool](#)[SAVANT](#)[SUAVE](#)[APS Documentation](#)[Contact Us](#)

Featured Stories


High-resolution (250m) baseline satellite bio-optical climatology developed for bays in northern Gulf of Mexico (CLICK HERE)

Tool developed to find and display current satellite imagery on a mobile device (See Mobile Image Tool link)

www7331.nrlssc.navy.mil/#

Bio-Optical/Physical Processes and Remote Sensing Section Code 7331 Stennis Space Center, Mississippi 39529

As a team, members of the Bio-Optical/Physical Processes and Remote Sensing Section have broad experience and expertise in many aspects of remote sensing and coupled physical/ecological modeling. We conduct research to better understand oceanographic processes in coastal and open-ocean environments. Our overarching goal is to exploit this knowledge of the marine environment to address a wide variety of navy needs related to optical variability, underwater light penetration, and physical/optical forecasting. However, we also work closely with external agencies and universities to address a wide variety of oceanographic science questions. Our state-of-the-art facilities include satellite receiving and image processing/analysis systems, in situ and laboratory optical instrumentation, advanced physical and ecological models, and access to high-performance super computers. Recent instrumentation acquisitions include gliders (one with an Optical Plankton Discriminator), trawl-resistant physical/optical moorings, and a flow cytometer. Research interests within the section include ocean color algorithm






United States Naval Research Laboratory
Bio-Optical/Physical Processes and Remote Sensing Section

[HOME](#)[BROWSE IMAGERY](#)[EXERCISE SUPPORT](#)[EXTERNAL LINKS](#)[PROJECTS](#)

- [Developing Ensemble Methods to Estimate Uncertainties in Remotely-Sensed Optical Properties \(DEMEN\)](#)
- [NRL 6.4 Project: Modeling, Sensing and Forecasting Ocean Optical Products for Navy Systems](#)
- [BIOSPACE \(Bio-optical Studies of Predictability and Assimilation in the Coastal Environment\)](#)
- [Dynamic Modeling of Marine Bioluminescence and Night Time Leaving Radiance](#)
- [Satellite Ocean Color Remote Sensing and Hydrodynamic Modeling to Understand Hypoxia in the Northern Gulf of Mexico](#)
- [Hyperspectral Imager for the Coastal Ocean \(HICO\)](#)
- [3D Remote Sensing with a Multiple-Band Active and Passive System: Theoretical Basis](#)
- [Characterizing Pelagic Habitats within U.S. Gulf of Mexico Coastal Waters Using Satellite Derived Data and Machine Learning Algorithms \(Eco-Mapping\)](#)
- [Application of a Global Aerosol Forecast Model for Multi-Spectral Ocean Color Atmospheric Corrections](#)
- [6.2 Resolving Bio-Optical Feedbacks to Ocean/Atmosphere Dynamics](#)
- [High-Resolution Subsurface Physical and Optical Property Fields in the Gulf of Mexico: Establishing Baselines and Assessment Tools for Resource Managers](#)
- [Automated Processing System \(APS\)](#)
- [Decadal-Scale Changes in Oceanic Heat Content for the Gulf of Mexico: A Model Study with Multi-disciplinary Implications to Climate Change](#)
- [Understanding and Forecasting Nonhydrostatic Dynamics in the Ocean](#)
- [6.1 Modeling Dynamic Bio-Optical Layers in Coastal Systems \(DYaBOLIC\)](#)
- [Transformational approach to monitoring water quality sustainability of coastal ecosystems from satellite remote sensing](#)
- [Development of Hyperspectral Imaging to Detect Contamination in Plants](#)
- [Improving Blended Multi-Sensor Ocean Color Products Through Assessment of Sensor Measurement Differences](#)
- [SAVANT \(SATellite VALidation NAVY Tool\)](#)
- [SUAVE \(Satellite Update Alerts for Various Exercises\)](#)

[Home](#)[Browse Imagery](#)[Exercise Support](#)[External Links](#)[Projects](#)[Help](#)[Seminars](#)[MISS BIGHT 2009
CHESAPEAKE BAY](#)[NRL-DC
NRL-SSC](#)[Current Projects](#)[Contact Webmaster](#)



High-Resolution Subsurface Physical and Optical Property Fields in the Gulf of Mexico: Establishing Baselines and Assessment Tools for Resource Managers

Objective: We propose to provide a database of subsurface physical and optical property fields for specific local areas within the Gulf of Mexico using a blend of data assimilative ocean modeling tools and ocean color remote sensing products. These three-dimensional property fields are needed to inform the decision-making process of resource managers, with particular emphasis on (i) habitat identification and characterization, (ii) assessment of the environmental impact of various permitted activities, such as dredging, and (iii) water quality assessment. Accordingly, we have identified two specific decision-making processes within the Gulf of Mexico that will be aided by the establishment of our proposed database. First, the Flower Garden Banks National Marine Sanctuary (FGBNMS) works with the Sanctuary Advisory Council, constituents, and users to conduct periodic management reviews of current regulations. The FGBNMS is currently going through a review, and potential boundary expansion has been identified as a priority management issue. The boundary expansion proposals and deliberations would be significantly aided by knowledge of baseline environmental conditions, such as light, temperature, and salinity, for benthic biological communities that may be ecologically linked to the presently protected areas. Second, the natural variability of coastal turbidity (suspended sediment load) is not known for local areas where the United States Army Corps of Engineers (USACE) are required to monitor during their dredging, flood protection, and coastal restoration activities. Knowledge of this natural variability would provide a baseline against which data from their monitoring activities may be assessed. With the expressed support of FGBNMS-NOAA and USACE, we will provide high-resolution and three-dimensional light, temperature, salinity, and current field estimates, and surface maps of suspended sediment load, made from a combination of satellite remote sensing and ocean modeling. By providing both a regional context and local-scale details, these products will help resource managers make more informed decisions.

[Image Climatology \(2005-2009\) Northern Gulf of Mexico Bays](#)






[Satellite-Extracted Data Files](#)

[Satellite Bio-Optics Project Summary Report](#)

Show me a Fancy list ▾ Sorted by Name ▾ Ascending ▾ in Normal order ▾ Matching * Go


NOTE: All SeaWiFS images and data presented on this website are for research and educational use only. All commercial use of SeaWiFS data must be coordinated with ORBIMAGE.

[Name](#) [Last modified](#) [Size](#) [Description](#)

-  [Parent Directory](#) -
-  [Choctawhatchee StAndrew Bays/](#) 14-Jan-2013 09:25 -
-  [Mobile Bay/](#) 14-Jan-2013 09:33 -
-  [Northern Gulf Coast/](#) 14-Jan-2013 09:41 -
-  [Pensacola Bay/](#)

[Name](#) [Last modified](#) [Size](#) [Description](#)

 [Parent Directory](#)

 [monthly time series/](#) 14-Jan-2013 09:25 -






[Name](#) [Last modified](#) [Size](#) [Description](#)

-  [Parent Directory](#) -
-  [Kd 488/](#) 14-Jan-2013 09:24 -
-  [PIM/](#) 14-Jan-2013 09:24 -
-  [POM/](#) 14-Jan-2013 09:24 -
-  [TSS/](#) 14-Jan-2013 09:25 -

[Name](#)

[Last modified](#) [Size](#) [Description](#)

 [Parent Directory](#)

-  [aqua.2005001.0101.2005031.0131.D.L4 MO.hmodis.PAN.v09.250m.Kd 488 lee.png](#) 04-Jan-2011 11:08 164K
-  [aqua.2005032.0201.2005059.0228.D.L4 MO.hmodis.PAN.v09.250m.Kd 488 lee.png](#) 04-Jan-2011 11:08 166K
-  [aqua.2005061.0301.2005091.0331.D.L4 MO.hmodis.PAN.v09.250m.Kd 488 lee.png](#) 08-Jan-2013 14:51 161K
-  [aqua.2005091.0401.2005120.0430.D.L4 MO.hmodis.PAN.v09.250m.Kd 488 lee.png](#) 04-Jan-2011 11:08 165K
-  [aqua.2005121.0501.2005151.0531.D.L4 MO.hmodis.PAN.v09.250m.Kd 488 lee.png](#) 04-Jan-2011 11:08 171K